

## CLONIDINE (tablet)

*klon-ih-deen*

### What does it do?

*Clonidine* is used to treat high blood pressure, severe pain, and sometimes other conditions.

### How should you take it?

Take *clonidine* tablets regularly as directed with a glass of water.

### What if you forget a dose?

Take the missed dose as soon as you remember. If it is close to the time for your next dose, skip the missed dose and carry on as normal. Do not take two doses at the same time.

### Can you take other medicines?

Some medicines available without a prescription may react with *clonidine* including:

- anti-inflammatories, such as diclofenac (e.g. Voltaren®), ibuprofen (e.g. Nurofen®), or aspirin (e.g. Disprin®, in doses used for pain relief). These can also be found in some cold and flu medicines (e.g. Nurofen Cold and Flu®).

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

### What side effects might you notice?

Side Effects	Recommended action
Slow heartbeat	Tell your doctor immediately
Drowsiness, dizziness, tiredness or weakness, headache Dry mouth, nausea, vomiting, constipation Low mood Changes in sexual function Joint, muscle or bone aches and pains	Tell your doctor if troublesome
Lightheaded or dizzy after standing up	Stand up slowly. If it continues, or is severe, tell your doctor

If you notice any other effects, discuss them with your doctor or pharmacist.

### Other information:

- Tell your doctor if you have heart or kidney problems, diabetes, or depression.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Do not stop taking *clonidine* without talking to your doctor as this may cause a sudden increase in your blood pressure.
- *Clonidine* may make you dizzy or sleepy and make it dangerous to drive, operate machinery or do other activities that require you to be alert. Limit alcohol intake because it can increase these effects.