



INSULIN LISPRO

in-sue-lin lis-pro

What does it do?

Insulin lispro is a rapid-acting insulin used to treat diabetes. It moves glucose from your blood into your cells, so the glucose can be used for energy.

Before you start

- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you use it?

Insulin lispro should be injected or infused under the skin. Change the injection or infusion site regularly so that you are less likely to get lumpiness under the skin.

Injection: The tummy is usually the best place to inject.

Infusion pump: The cannula can be placed on your tummy, upper arm, hips, thighs, or lower back.

Do not stop using *insulin lispro* without talking to your health professional first.

What if you forget a dose?

If you miss a dose of *insulin lispro* and have already eaten, check your blood glucose. If it is high, follow your instructions for high blood glucose (hyperglycemia). Otherwise, skip the dose and continue as directed. If you are unsure, contact your health professional.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|--|---|
| Low blood glucose (hypo): symptoms may include sweating, trembling, feeling anxious or irritable | Drink or eat something sweet. Tell your health professional if this happens a lot or is severe. |
| Weight gain Pain, tenderness or redness at injection site | Tell your health professional if troublesome |
| Lumpiness under the skin | Change the injection or infusion site regularly. Discuss with your health professional. |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Check your blood glucose levels before meals, 2 hours after meals and at bedtime, or as directed.
- Wear medical identification (e.g. MedicAlert bracelet) saying that you have diabetes. Always keep insulin and something sweet with you.
- You will need a regular blood test (HbA1c) to check how your diabetes is controlled.
- If you are unwell, follow your [sick-day plan](#), as the amount of insulin you need may change. Discuss this with your health professional.
- Exercise or physical activity may change the amount of insulin you need. Discuss this with your health professional.
- Alcohol may change your blood glucose levels and make you more likely to have a hypo. It may also mask your warning signs of low blood glucose. See [Alcohol and Diabetes](#).
- Check your blood glucose levels before driving. Do not drive if you are having a hypo as low blood glucose could make driving unsafe. See [Driving with diabetes](#)
- Keep unopened insulin in the fridge. Once you start using it, you can keep it at room temperature for about 4 weeks (check the exact time on the packet for your insulin). After this, take any leftover insulin back to your pharmacy.

This leaflet contains important, but not all, information about this medicine.

Prepared by the MyMedicines Committee at Christchurch Hospital, Te Whatu Ora - Waitaha, New Zealand. November 2024