



PETHIDINE

peth-ih-deen

What does it do?

Pethidine is used to relieve strong pain.

How should you take it?

Take *pethidine* as directed with a glass of water.

What if you forget a dose?

If *pethidine* is taken regularly, take the missed dose as soon as possible. If it is close to the time for your next dose, skip the missed dose and carry on as normal. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *pethidine* including:

- sedating antihistamines (e.g. Phenergan®)
- cough suppressants (e.g. Duro-Tuss®, Benadryl Dry Forte®)
- pain relief medicines containing codeine (e.g. Panadeine®)
- some migraine medicines, such as sumatriptan (e.g. Sumagran Active®) or zolmitriptan (e.g. Zomig®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. valerian) or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Slow or shallow breathing, hard to wake up Change in heartbeat Seizures	Tell your doctor immediately
Constipation Confusion, dizziness, agitation, hallucinations, mood swings	Tell your doctor
Drowsiness, weakness, headache, trouble sleeping Nausea, vomiting, stomach upset Dry mouth or throat, changes in vision, trouble peeing Swollen feet or legs Sweating	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have lung, bowel, liver, kidney, bladder, pancreas or heart problems, if you have ever had a seizure, or have had a recent head injury.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- If *pethidine* is not relieving your pain contact your health professional – your dose may need changing.
- *Pethidine* may make you dizzy or sleepy and make it dangerous to drive, operate machinery or do other activities that require you to be alert. Limit alcohol intake because it can increase these effects.
- *Pethidine* may be addictive with long-term use.