



IPRATROPIUM AND SALBUTAMOL

eye-pra-troe-pee-um and sal-bew-ta-mol

What does it do?

Ipratropium and *salbutamol* opens narrowed airways in the lungs. It helps to relieve asthma symptoms and breathing problems.

How should you use it?

Inhale *ipratropium* and *salbutamol* as directed by your healthcare professional.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and inhale your next dose at the usual time. Otherwise, inhale the missed dose as soon as you remember. Do not inhale two doses at the same time.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|---|---------------------------------|
| Eye pain, changes in vision | Tell your doctor immediately |
| Fast or irregular heartbeat, chest pain | Tell your doctor |
| Tremor, muscle cramps Headache, dizziness Nervousness, trouble sleeping Dry mouth, trouble peeing Hoarse voice, unpleasant taste, cough | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have heart, bladder or prostate problems, high blood pressure or glaucoma.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Take care not to get the spray from the *ipratropium* and *salbutamol* inhaler or nebuliser into your eyes. If this happens, rinse your eyes thoroughly with water.
- Some people with asthma may get short of breath after taking certain pain-relieving medicines such as aspirin (e.g. *Aspec*®), diclofenac (e.g. *Voltaren*®) or ibuprofen (e.g. *Nurofen*®). Talk to your healthcare professional.
- The use of inhalers can sometimes make breathing worse. Discuss this with your healthcare professional.