



DISOPYRAMIDE

dye-so-pir-ah-mide

What does it do?

Disopyramide is used to treat and prevent irregular heartbeats.

How should you take it?

Take *disopyramide* regularly as directed with a glass of water. You can take it with or without food.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *disopyramide* including:

- sildenafil (e.g. Viagra®) or similar medicines
- some antihistamines or anti-nausea medicines such as promethazine (e.g. Phenergan®), hyoscine (e.g. Scopoderm®), meclozine (e.g. Sea-legs®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|---|---------------------------------|
| Symptoms of heart problems including: fluttering, pounding or pain in your chest, feeling your heart skips a beat, dizziness or fainting | Tell your doctor immediately |
| Low blood sugar: symptoms may include sweating, trembling, feeling anxious or irritable Swollen feet or legs, short of breath Confusion | Tell your doctor |
| Dry mouth, blurred vision, constipation or trouble peeing (anticholinergic effects) Nausea | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have kidney, prostate, bladder or bowel problems, diabetes or angle-closure glaucoma.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- *Disopyramide* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)