



PRIMIDONE

prim-ih-dohn

What does it do?

Primidone is used to treat and prevent seizures. It is also sometimes used for other conditions.

Before you start

- Tell your doctor if you have liver problems or sleep apnoea.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *primidone* regularly as directed with a glass of water. You can take it with or without food.

What if you forget a dose?

Take the missed dose as soon as possible and continue as directed.

Can you take other medicines?

Primidone can react with many medicines, sometimes with severe results.

Remember to tell your pharmacist or doctor you are taking *primidone* before starting any other medicines or treatments, including over-the-counter medicines, vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|---|---|
| Skin rash, skin peeling or blisters | Stop taking and see your doctor immediately |
| Suicidal thoughts Slow or shallow breathing, hard to wake up Reduced number of blood cells that fight infections or help your blood to clot - symptoms include: fever, chills, sore throat or generally feeling unwell, or easy or unusual bruising or bleeding Symptoms of liver problems including: yellow skin or eyes, itching, dark urine, pale bowel motions, abdominal pain | Tell your doctor immediately |
| Loss of co-ordination/walking or handwriting problems Confusion, trouble concentrating Unusual behaviour or thinking, mood changes, agitation | Tell your doctor |
| Dizziness, drowsiness | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Some contraceptives may not work as well while you are taking *primidone*, and for 4 weeks after stopping. Discuss with your doctor or pharmacist.
- *Primidone* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- Long-term use of *primidone* may weaken your bones.
- Do not stop taking *primidone* without talking to your doctor first, unless you have a skin rash (see Side Effects).