



ROXITHROMYCIN

rox-ith-roe-my-sin

What does it do?

Roxithromycin is used to treat and prevent bacterial infections.

How should you take it?

Take *roxithromycin* regularly as directed with a glass of water. Keep taking it until the course is finished, even if you start to feel better.

What if you forget a dose?

Take the missed dose as soon as possible and continue as directed.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Fast or irregular heartbeat, fainting Symptoms of liver problems including: yellow skin or eyes, itching, dark urine, pale bowel motions, abdominal pain	Tell your doctor immediately
Severe or persistent diarrhoea, abdominal pain Small white sores in mouth, furry tongue (oral thrush) Vaginal itch or discharge (vaginal thrush)	Tell your doctor
Diarrhoea Headache	Tell your doctor if troublesome
Stomach upset	Take with food

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have heart, liver or gut problems.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Your doctor may do a heart test (ECG) before you start and while you are taking *roxithromycin*.