

RIZATRIPTAN

ri-zah-trip-tan

What does it do?

Rizatriptan is used to treat migraines.

Before you start

- Tell your health professional if you have heart or blood vessel problems, high blood pressure, or if you have ever had a stroke, or 'mini-stroke'.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Tell your doctor if you have phenylketonuria (PKU). *Rizatriptan* wafers contain aspartame – a source of phenylalanine.

How should you take it?

Take *rizatriptan* as directed when your symptoms first start. Place it on your tongue and allow it to dissolve, then swallow.

Do not remove the wafer blister from the outer sachet until just before taking the dose. The blister pack should be peeled open with clean, dry hands.

If your symptoms get better, but then return, you can take another dose after 2 hours.

If *rizatriptan* does not help, do not take any further doses for the same attack, as it is unlikely to work. You can still use *rizatriptan* for a new attack.

Do not take more than the maximum daily dose advised by your doctor.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Pain, tightness or pressure in your chest, throat or jaw Fast or irregular heartbeat	Tell your doctor immediately
Drowsiness, tiredness or weakness, dizziness Hot flushing Burning sensation, tingling or numbness	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you need to use *rizatriptan* several times each week. Using *rizatriptan* too often can cause a different type of headache. Your doctor can talk with you about different medicines that are taken every day to prevent migraines.
- Do not use *rizatriptan* to prevent migraines.
- *Rizatriptan* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)