

ALLOPURINOL

al-oh-pure-ih-nol

What does it do?

Allopurinol is used to prevent gout and kidney stones. It reduces the amount of uric acid made in your body.

How should you take it?

Take *allopurinol* regularly as directed. Take with food and a glass of water.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Skin rash, itching, fever	Stop taking and see your doctor immediately
Trouble peeing Gout attack Headache, dizziness, drowsiness	Tell your doctor if troublesome
Stomach upset	Take with food

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Tell your doctor if you have kidney or liver problems.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- *Allopurinol* can cause kidney stones if you do not drink enough fluid. Discuss with your healthcare professional.
- *Allopurinol* can cause a gout attack when you start taking it, so your doctor may prescribe another medicine to prevent this happening. Keep taking *allopurinol* during a gout attack. Stopping the tablets suddenly is likely to make your gout worse.