

## CEFACTOR

*kef-ah-klor*

### What does it do?

*Cefactor* is used to treat and prevent bacterial infections.

### How should you take it?

Take *cefactor* regularly as directed. Keep taking it until the course is finished, even if you start to feel better.

Take the capsules with a glass of water.

Measure the liquid carefully with an oral syringe or measuring spoon. Shake the bottle well before measuring each dose.

### What if you forget a dose?

Take the missed dose as soon as possible and continue as directed.

### Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

### What side effects might you notice?

Side Effects	Recommended action
Symptoms of allergy including: skin rash, itching, swelling, trouble breathing	Tell your doctor immediately
Joint, muscle or bone aches and pains, fever Severe or persistent diarrhoea, abdominal pain Small white sores in mouth, furry tongue (oral thrush) Vaginal itch or discharge (vaginal thrush)	Tell your doctor
Diarrhoea, stomach upset	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

### Other information:

- Tell your doctor if you have kidney or gut problems, have ever had a seizure, or have ever had an allergic reaction to cephalosporins, penicillins or any other medicine.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Store *cefactor* liquid as directed. Once the course is finished, take any leftover liquid back to your pharmacy.