New Zealand Formulary
PATIENT INFORMATION

# **COVID-19 VACCINE (Pfizer - BioNTech)**

## What does it do?

The COVID-19 vaccine helps protect you against severe COVID-19 infection. It means you could have milder or no COVID-19 symptoms and recover faster.

# Before you start

- Tell your health professional if you currently feel unwell or have a fever.
- Tell your health professional if you have HIV, receive dialysis, take medicines which weaken your immune system, or if you have had heart inflammation (myocarditis) after a previous COVID-19 vaccine.
- Tell your health professional if you are pregnant, planning to become pregnant, or breastfeeding.

# How is it given?

The COVID-19 vaccine is given as an injection into a muscle. It will be given to you by a health professional.

You will need to wait for at least 15 minutes after having your vaccine, to check you don't have an allergic reaction.

#### What if you forget a dose?

You can manage your booking online at bookmyvaccine.covid19.health.nz or by phoning 0800 28 29 26.

#### Can you take other medicines?

Tell your pharmacist or doctor about <u>all medicines or treatments</u> that you may be taking, including vitamins, herbal products or recreational drugs.

## What side effects might you notice?

Side Effects	Recommended action
Flu-like symptoms (fever, chills, aches and pains, nausea, headache, tiredness or weakness) Pain, tenderness or redness at injection site	This usually passes after a few days, take paracetamol if needed. Tell your health professional if troublesome.
Symptoms of allergy including: skin rash, itching, swelling, trouble breathing	Tell your health professional immediately
Symptoms of heart inflammation including: fluttering, pounding or pain in your chest, shortness of breath	Tell your doctor immediately

If you notice any other effects, discuss them with your doctor or pharmacist.

#### Other information:

• The COVID-19 vaccine may cause swelling of your lymph nodes (a lump in your armpit or neck). Tell your doctor if they are still swollen after 6 weeks. If you have a CT scan or breast screening appointment, tell your health professional you have had your vaccine within the last 6 weeks.