

## CYCLIZINE

*syé-kli-zeen*

### What does it do?

*Cyclizine* is an antihistamine used to treat and prevent nausea and vomiting.

### Before you start

- Tell your doctor if you have bladder, prostate or bowel problems, or angle-closure glaucoma.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

### How should you take it?

Take *cyclizine* as directed with a glass of water. You can take it with or without food.

### What if you forget a dose?

If you take *cyclizine* regularly and it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

### Can you take other medicines?

Some medicines available without a prescription may react with *cyclizine* including:

- some antihistamines or anti-nausea medicines such as promethazine (e.g. Phenergan®), hyoscine (e.g. Scopoderm®), meclozine (e.g. Sea-legs®)
- other medicines that may cause drowsiness (e.g. pain-killers, anti-nausea and cough/cold medicines – ask your pharmacist)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

### What side effects might you notice?

Side Effects	Recommended action
Confusion	Tell your doctor
Dry mouth, blurred vision, constipation or trouble peeing (anticholinergic effects) Drowsiness	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

### Other information:

- *Cyclizine* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)