



ETIDRONATE

et-ih-dron-ate

What does it do?

Etidronate is used to treat osteoporosis (weak bones). It helps to make your bones stronger and less likely to break. It is also sometimes used for other conditions.

Before you start

- Tell your doctor if you have kidney, parathyroid, dental or swallowing problems, or if you get heartburn or indigestion.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *etidronate* as directed. You need to take it at least 2 hours before or after food or drink and other medicines. *Etidronate* can damage your oesophagus (food pipe). To avoid this, take it with a large glass of water. Swallow the tablet whole, do not crush or chew it. Sit or stand upright for at least 30 minutes after taking a dose.

Etidronate is usually taken for 2 weeks every 3 months.

What if you forget a dose?

Skip the missed dose and take the next day's dose at the usual time.

Can you take other medicines?

Some medicines available without a prescription may react with *etidronate* including:

- calcium supplements (e.g. Calci-Tab®)
- iron supplements (e.g. Ferro-Tab®)
- antacids (e.g. Mylanta®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Trouble swallowing, chest pain Indigestion or heartburn (new or getting worse) Eye pain, changes in vision	Tell your doctor immediately
Joint, muscle or bone aches and pains Stomach upset	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Before you have any dental work, tell your dentist you are taking *etidronate*.