



MEDROXYPROGESTERONE DEPOT (injection)

me-drox-ee-pro-jes-te-rone dee-poe

What does it do?

Medroxyprogesterone depot is used to prevent pregnancy. It may also be used to treat endometriosis and some cancers.

Before you start

- Tell your doctor if you have breast cancer.

How is it given?

Medroxyprogesterone depot is given as an injection into a muscle by your health professional.

To prevent pregnancy, *medroxyprogesterone depot* is usually given every 12 weeks.

You can start at any time in your cycle. If you start within 5 days of the first day of your period, you are protected from pregnancy straight away. If you start more than 5 days after the first day of your period, you will need to use condoms (or don't have sex) for the next 7 days.

What if you forget a dose?

If you miss an appointment for your injection, contact your health professional as soon as possible.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|--|--|
| Changes in periods | Irregular bleeding is common in the first 3-6 months. This is not harmful and usually improves with time. After this, some women will not have any periods at all. |
| Mood changes, headache Acne Weight gain Sore breasts Stomach upset | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Medroxyprogesterone depot* may weaken your bones, especially if you are using it for longer than two years. This usually improves after you stop using it.
- *Medroxyprogesterone depot* does not protect you from sexually transmitted infections.
- If you want to get pregnant after stopping *medroxyprogesterone depot*, you may find that it can take up to a year.