

FLUCLOXACILLIN (injection)

floo-klox-ah-sil-in

What does it do?

Fluctorial infections and prevent bacterial infections.

Before you start

- Tell your doctor if you have kidney or liver problems, or if you are allergic to penicillin antibiotics.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How is it given?

Flucloxacillin is given as an infusion or injection into a vein, or as an injection into a muscle. It is usually given to you by a health professional.

What if you forget a dose?

Contact your health professional as soon as possible. Usually, the missed dose is given as soon as possible, and treatment continued as directed.

Can you take other medicines?

Tell your pharmacist or doctor about <u>all medicines or treatments</u> that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|---|--|
| Symptoms of allergy including: skin rash, itching, swelling, trouble breathing Symptoms of liver problems including: yellow skin or eyes, itching, dark urine, pale bowel motions, abdominal pain | Tell your health professional immediately |
| Severe or persistent diarrhoea, abdominal pain Small white sores in mouth, furry tongue (oral thrush) Vaginal itch or discharge (vaginal thrush) | Tell your health professional |
| Stomach upset Imitation or pain at injection site | Tell your health professional if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.