

EMTRICITABINE

emtry-si-tab-een

What does it do?

Entricitabine is used with other medicines to control HIV.

Before you start

- Tell your doctor if you have kidney problems, or if you have ever had Hepatitis B or C.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *entricitabine* as directed with a glass of water. It is very important to take it regularly. If you often forget to take *entricitabine*, it may not work as well.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *entricitabine* including:

orlistat (Xenical®)

Tell your pharmacist or doctor about <u>all medicines or treatments</u> that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Headache, dizziness Trouble sleeping, strange dreams Joint, muscle or bone aches and pains Cough, runy or blocked nose Darkening of skin or fingemails Stomach upset, abdominal pain	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- You will need regular blood tests while taking entricitabine to check your viral load.
- Entricitabine belongs to a group of medicines that have caused a serious condition called lactic acidosis.
 Contact your doctor immediately if you suddenly feel very unwell or unusually tired, or if your breathing becomes faster than normal.
- As your HIV comes under control, your immune system begins to recover. It may start fighting infections better
 than before which may make you unwell for a while. Your doctor may call this Immune Reconstitution
 Syndrome.
- Treatment for HIV may change your body shape. These changes can include fat build-up, fat loss, or both.
 Discuss this with your doctor.
- Do not stop taking entricitabine without talking to your doctor first.