Te Whatu Ora
Health New Zealand
Waitaha Canterbury

PRETOMANID

pre-toe-ma-nid

What does it do?

Pretomanid is an antibiotic used with other medicines to treat tuberculosis (TB).

Before you start

- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Your doctor may do a heart test (ECG) before you start and while you are taking *pretomanid*.

How should you take it?

Take *pretomanid* regularly as directed with food and a glass of water. Keep taking it until the course is finished.
It is important to take it regularly. If you often forget to take it, it may not work as well.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Fainting (may be a heartbeat problem)	Tell your doctor immediately
Stomach upset	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Pretomanid* is not registered for use in New Zealand. Discuss with your doctor.