

INSULIN DEGLUDEC

in-sue-lin deg-loo-dek

What does it do?

Insulin degludec is a long-acting insulin used to treat diabetes. It moves glucose from your blood into your cells, so the glucose can be used for energy.

Before you start

- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you use it?

Insulin degludec should be injected under the skin. The tummy is usually the best place to inject. Change the injection site regularly so that you are less likely to get lumpiness under the skin.

Do not stop using *insulin degludec* without talking to your health professional first.

What if you forget a dose?

If it is almost time for your next dose of *insulin degludec*, skip the dose you missed and continue at your normal time. Do not inject two doses at the same time. Otherwise, inject the missed dose as soon as possible. If you are unsure, contact your health professional.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|--|---|
| Low blood glucose (hypo): symptoms may include sweating, trembling, feeling anxious or irritable | Drink or eat something sweet. Tell your health professional if this happens a lot or is severe. |
| Weight gain Pain, tenderness or redness at injection site | Tell your health professional if troublesome |
| Lumpiness under the skin | Change the injection site regularly. Discuss with your health professional. |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- Check your blood glucose levels before meals, 2 hours after meals and at bedtime, or as directed.
- Wear medical identification (e.g. MedicAlert bracelet) saying that you have diabetes. Always keep insulin and something sweet with you.
- You will need a regular blood test (HbA1c) to check how your diabetes is controlled.
- If you are unwell, follow your sick-day plan as the amount of insulin you need may change. Discuss this with your health professional.
- Exercise or physical activity may change the amount of insulin you need. Discuss this with your health professional.
- Alcohol may change your blood glucose levels and make you more likely to have a hypo. It may also mask your warning signs of low blood glucose.
- Check your blood glucose levels before driving. Do not drive if you are having a hypo as low blood glucose could make driving unsafe.
- Keep unopened insulin in the fridge. Once you start using it, you can keep it at room temperature for about 4 weeks (check the exact time on the packet for your insulin). After this, take any leftover insulin back to your pharmacy.
- *Insulin degludec* is not registered for use in New Zealand. Discuss with your doctor.

This leaflet contains important, but not all, information about this medicine.

Prepared by the MyMedicines Committee at Christchurch Hospital, Te Whatu Ora - Waitaha, New Zealand. July 2024