

TAMSULOSIN

tam-soo-low-sin

What does it do?

Tamsulosin is used to treat trouble peeing caused by an enlarged prostate.

Before you start

- Tell your doctor if you have heart problems, low blood pressure, or problems with fainting.

How should you take it?

Take *tamsulosin* regularly as directed with a glass of water. Swallow the capsules whole. You can take it with or without food.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *tamsulosin* including:

- sildenafil (e.g. Viagra®) or similar medicines

Tell your pharmacist or doctor about [all medicines or treatments](#) that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|--|--|
| Tiredness or weakness Runny or blocked nose Dry orgasm | Tell your doctor if troublesome |
| Lightheaded or dizzy after standing up | Stand up slowly. If it continues, or is severe, tell your doctor |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Tamsulosin* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- If you are having eye surgery (e.g. cataract operation), tell your doctor that you are taking *tamsulosin*.