

BENZATROPINE

ben-zah-troe-peen

What does it do?

Benzatropine is used to treat Parkinson's disease. It balances levels of chemicals in the brain to help with symptoms such as movement problems. It is also used for other conditions.

Before you start

- Tell your doctor if you have bladder, prostate, bowel or mental health problems, or angle-closure glaucoma.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *benzatropine* regularly as directed with a glass of water. You can take it with or without food.

Do not stop taking *benzatropine* suddenly without talking to your doctor first. Your doctor may reduce the dose gradually.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *benzatropine* including:

- some antihistamines or anti-nausea medicines such as promethazine (e.g. Phenergan®), hyoscine (e.g. Scopoderm®), meclozine (e.g. Sea-legs®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

| Side Effects | Recommended action |
|------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------|
| Trouble thinking clearly or concentrating Unusual behaviour or thinking Seeing, hearing or feeling things that aren't really there | Tell your doctor |
| Dry mouth, blurred vision, constipation or trouble peeing (anticholinergic effects) Stomach upset | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Benzatropine* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. See [Are you safe to drive?](#)