

PRAMIPEXOLE

pram-e-pex-ole

What does it do?

Pramipexole is used to treat Parkinson's disease and sometimes other conditions. It acts like a chemical in your brain called dopamine.

Before you start

- Tell your doctor if you have kidney or mental health problems, or low blood pressure.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *pramipexole* regularly as directed with a glass of water.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Hallucinations, confusion, unusual behaviour or thinking Unusual urges (e.g. gambling, eating, spending, sex) Swollen feet or legs, short of breath Falling asleep without warning	Tell your doctor
Drowsiness Dizziness, headache Strange or uncontrolled movements Nausea, constipation	Tell your doctor if troublesome
Lightheaded or dizzy after standing up	Stand up slowly. If it continues, or is severe, tell your doctor

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Pramipexole* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- If you are taking *pramipexole* for restless leg syndrome, your symptoms might get worse when you start. Discuss this with your doctor, as changing the dose can help.
- Do not stop taking *pramipexole* without talking to your doctor first.