

CITALOPRAM

sit-al-oh-pram

What does it do?

Citalopram is used to treat depression, and sometimes other conditions.

Before you start

- Tell your doctor if you have bleeding problems, bipolar disorder, or if you have ever had a seizure.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Your doctor may do a heart test (ECG) before you start and while you are taking *citalopram*.

How should you take it?

Take *citalopram* regularly as directed with a glass of water. You can take it with or without food.

What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

Can you take other medicines?

Some medicines available without a prescription may react with *citalopram* including:

- anti-inflammatories, such as diclofenac (e.g. Voltaren®), ibuprofen (e.g. Nurofen®), or aspirin (e.g. Disprin®, in doses used for pain relief). These can also be found in some cold and flu medicines (e.g. Nurofen Cold and Flu®).
- low-dose aspirin (e.g. Cartia®)
- some migraine medicines, such as sumatriptan (e.g. Sumagran Active®)
- omeprazole (e.g. Losec®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs (e.g. ecstasy).

What side effects might you notice?

| Side Effects | Recommended action |
|--|---------------------------------|
| Suicidal thoughts Fainting Muscle twitching or shaking you can't control, confusion, heavy sweating, fever Seizures | Tell your doctor immediately |
| Anxiety, restlessness Easy or unusual bruising or bleeding | Tell your doctor |
| Nausea, vomiting, diarrhoea, loss of appetite, dry mouth Drowsiness, trouble sleeping, tiredness or weakness, yawning Dizziness, headache, sweating Less interest in, or trouble having sex | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- *Citalopram* can impair your ability to do tasks such as driving or using machines. Alcohol makes this worse. Discuss your risk with your health professional. (search NZTA - Are you safe to drive?)
- It may take a few weeks for *citalopram* to start working.
- Do not stop taking *citalopram* without talking to your doctor first.