

WARFARIN

war-far-in

What does it do?

Warfarin is used to treat and help prevent clots in your blood. It reduces your chance of having a stroke and new clots forming in your legs or lungs. It is also used to help prevent clots in people who have had a mechanical heart valve replacement.

Before you start

- Tell your doctor if you have bleeding problems.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.

How should you take it?

Take *warfarin* regularly as directed with a glass of water. Take it at the same time each day.

Your dose will depend on a blood test called an INR. This measures how long it takes your blood to clot. Your health professional will tell you what your INR is and what dose to take. It is important to record your INR and dose.

Warfarin comes in two brands in NZ (Marevan® and Coumadin®). Check which brand you have and stick to it.

Warfarin comes in different strengths, with different colours. Most people take the Marevan® brand shown in the picture below. If your tablets look different than usual, check with your pharmacist.

Marevan® tablets



What if you forget a dose?

Take the missed dose if you remember on the same day. If not, skip the dose and carry on as normal. Do not take two doses on the same day. Record any missed doses and tell your health professional on your next visit.

Can you take other medicines?

Some medicines available without a prescription may react with *warfarin* including:

- anti-inflammatories, such as diclofenac (e.g. Voltaren®), ibuprofen (e.g. Nurofen®), or aspirin (e.g. Disprin®, in doses used for pain relief). These can also be found in some cold and flu medicines (e.g. Nurofen Cold and Flu®).
- low-dose aspirin (e.g. Cartia®) - sometimes prescribed with *warfarin*, check with your doctor.

Warfarin can react with many medicines, sometimes with severe results. Remember to tell your pharmacist or doctor you are taking *warfarin* before starting any other medicines or treatments, including over-the-counter medicines, vitamins, herbal products or recreational drugs.

What side effects might you notice?

Side Effects	Recommended action
Bleeding you cannot stop, bruises that get bigger, coughing or vomiting blood, vomit that looks like coffee grounds, red or dark brown urine, red or black poo, severe headache (may be signs of bleeding) Skin rash, change in skin colour (blue or purple)	Tell your doctor immediately
Hair loss or thinning	Tell your doctor if troublesome

If you notice any other effects, discuss them with your doctor or pharmacist.

Other information:

- It is important to tell anyone who gives you medical or dental treatment (e.g. doctor, dentist, pharmacist or podiatrist) that you are taking *warfarin*.
- You may bleed or bruise more easily while you are taking *warfarin*. Take care or avoid activities that may cause bleeding e.g. tattoos, piercings and contact sports.
- If you have a serious fall or hurt your head, talk to or see a doctor straight away.
- Changes in your diet may affect your INR. Talk to your health professional.
- Alcohol can increase your chance of bleeding. You can still have the occasional drink, but avoid binge drinking.