

## TADALAFIL

*tah-dah-lah-fil*

### What does it do?

*Tadalafil* is used to treat pulmonary hypertension (high blood pressure in the blood vessels of your lungs), and sometimes other conditions. It improves blood flow by relaxing your blood vessels.

If you are taking *tadalafil* for erectile dysfunction see separate information sheet "*Tadalafil* (for erectile dysfunction)".

### Before you start

- Tell your doctor if you have had a recent stroke, heart or blood pressure problems, or an eye problem called optic neuropathy.
- Tell your doctor if you are pregnant, planning to become pregnant, or breastfeeding.
- Do not use *tadalafil* if you are taking a nitrate medicine (e.g. glyceryl trinitrate or isosorbide mononitrate). Taking them both can cause a dangerous drop in blood pressure.

### How should you take it?

Take *tadalafil* regularly as directed with a glass of water. You can take it with or without food.

### What if you forget a dose?

If it is nearly time for your next dose, skip the missed dose and take your next dose at the usual time. Otherwise, take the missed dose as soon as you remember. Do not take two doses at the same time.

### Can you take other medicines?

Some medicines available without a prescription may react with *tadalafil* including:

- glyceryl trinitrate (GTN) (e.g. Nitrolingual®, Glytrin®)

Tell your pharmacist or doctor about all medicines or treatments that you may be taking, including vitamins, herbal products (e.g. St John's wort) or recreational drugs.

### What side effects might you notice?

| Side Effects  | Recommended action              |
|---|---------------------------------|
| Prolonged erection (longer than four hours)<br>Changes in vision                                  | Tell your doctor immediately    |
| Headache, dizziness<br>Flushing<br>Runny or blocked nose<br>Muscle aches and pains<br>Indigestion | Tell your doctor if troublesome |

If you notice any other effects, discuss them with your doctor or pharmacist.

### Other information:

- Grapefruit, grapefruit juice or sour/Seville oranges may react with *tadalafil*. Discuss with your pharmacist.
- Do not stop taking *tadalafil* without talking to your doctor first.